The Science Case for Simultaneous mm-Wavelength Receivers in Radio Astronomy

Editor: María J. Rioja

Contributors: Richard Dodson, Taehyun Jung, José Luis Goméz, Valentin Bujarrabal, Luca Moscadilli, James C. A. Miller-Jones, Alexandra J. Tetarenko, Gregory R. Sivakoff

Informe Técnico IT - CDT&OAN 2016 - 24

Abstract

Presentamos un resumen de las aplicaciones científicas discutidas en el Foro Técnico Europeo de Radioastronomía (ERATec) celebrado en Florencia, en Octubre de 2015, para resaltar el amplio hábito de aplicación y el impacto derivados del uso de receptores con capacidad para observaciones simultáneas a múltiples frecuencias en el rango de longitudes de onda milimétricas.

Este informe pretende impulsar la aplicación de las nuevas tecnologías y métodos, que ya están disponibles, en los radio telescopios europeos para avanzar los estudios en una amplia gama de campos de investigación en astronomía.

Describimos las nuevas oportunidades y ventajas que ofrecen para el estudio en los siguientes campos: observaciones con interferometría de muy larga base (VLBI) de la emisión en radio continuo (estudios de objetos celestes débiles, medidas de astrometría de alta precisión, estudios de núcleos activos de galaxias (AGNs) con medidas de índices espectrales y rotaciń de Faraday), observaciones con VLBI de la emisión por medio de líneas espectrales (estudios de regiones de formación de estrellas evolucionadas y masivas) y medidas de las variaciones en el dominio temporal de la emisión en radio frecuencias procedente de chorros compactos en sistemas binarios con emisión de rayos X.

Nuestra recopilación reúne una amplia gama de importantes aplicaciones científicas que se beneficiarán enormemente de la capacidad de observaciones simultáneas a múltiples frecuencias en el rango de longitudes de onda milimétricas.

Instrumentos con dicha capacidad son esenciales para mejorar la eficiencia, sensibilidad, precisión y solidez de las aplicaciones científicas mencionadas. En algunos casos, sin receptores simultáneos los objetivos científicos son simplemente inalcanzables.

Por lo tanto, alentamos firmemente a los observatorios europeos para que tomen la delantera en la instalación de tales capacidades en sus instalaciones.

Contents

1	Int	roduc	tion	3			
2	Continuum Studies						
	2.1	Weak	Sources	4			
	2.2	Astro	metry	7			
	2.3	Faraday Rotation studies					
	2.4	Opacity core-shifts, γ -ray flares, and the nature of the VLBI core 12					
3	Spectral Studies						
	3.1	Multifrequency VLBI observations of maser emission in evolved stars					
		3.1.1	AGB and post-AGB evolutionary phases	13			
		3.1.2	SiO and H_2O masers from AGB stars $\ldots \ldots \ldots \ldots \ldots \ldots \ldots$	14			
		3.1.3	Maser lines in proto-PNe	17			
		3.1.4	Conclusions	18			
	3.2 Massive Stars						
		3.2.1	$\mathbf{Introduction}\ .\ .\ .\ .\ .\ .\ .\ .\ .\ .\ .\ .\ .\$	19			
		3.2.2	Source I in Orion BN/KL	20			
		3.2.3	Advantages of multi-frequency maser observations	21			
		3.2.4	Prospects for the Future	22			
4	Tir	ne do	omain observations of X-ray binary jets	23			
5	5 Concluding Remarks						

1. Introduction

Very Long Baseline Interferometry (VLBI) studies at cm wavelengths is a well-established field, with advanced technological developments and analysis techniques that result in superb quality images, including those of very weak μ Jy sources (e.g., Garrett 2005) and with microarcsecond (μ as) astrometry (e.g., Reid & Honma 2014), using phase referencing (*hereafter* PR) techniques.

VLBI at mm and sub-mm wavelengths (*hereafter* mm-VLBI) can result in the highest angular resolutions achieved in astronomy and has a unique access to emission regions that are inaccessible with any other approach or at longer wavelengths, because the compact areas of interest are often self-absorbed or scatter-broadened. Therefore it holds the potential to increase our understanding of the physical processes in, for example, Active Galactic Nuclei (AGN) and in the vicinity of super-massive black holes, and for studies of molecular transitions at high frequencies.

Nevertheless the scientific applications of mm-VLBI have to date been much less widespread. The reason being that the observations become progressively more challenging as the wavelength gets shorter because of the: limited telescope surface accuracy and efficiency, higher receiver system temperatures and lower sensitivity, shorter atmospheric coherence times and the fact that sources are intrinsically weaker in general. These in turn prevent the use of phase referencing calibration techniques, which are routinely used in cm-VLBI, and all benefits resulting from them, beyond \sim 43 GHz (with a single exception of PR at 86 GHz with the VLBA by Porcas & Rioja 2002).

Continuous development and technical improvements have led to a sustained increase of the high frequency threshold for VLBI observations in the last two decades (e.g., Krichbaum et al. 2014). Regular observations up to 86 GHz are being carried out with well established networks such as the Very Long Baseline Array (VLBA) and Global mm-VLBI Array (GMVA), most recently with the Korean VLBI Network (KVN) up to 130 GHz, and ad-hoc observations at the highest frequencies up to 240 GHz with the Event Horizon Telescope (EHT) (Doeleman et al. 2008). The field of mm-VLBI will greatly benefit from the arrival of phased-up Atacama Large Millimeter and submillimeter Array (ALMA) (Matthews & Crew 2015) for joint VLBI observations.

The benefit of multi-frequency based calibration techniques in observations at high frequencies, which are dominated by non-dispersive fast tropospheric fluctuation errors, has long been known. It relies on using the correctly scaled calibration derived at lower frequencies to correct the higher frequencies; see, for example, Carilli & Holdaway (1999); Asaki et al. (1996) for connected interferometers. This has been extended to VLBI observations using the frequency agility capability (i.e. fast frequency switching) of the VLBA, initially offering an increased coherence (Middelberg et al. 2005) and, after additional development, bona-fide astrometric measurements (Dodson & Rioja 2009; Rioja & Dodson 2011) at the highest frequency of 86 GHz. The use of this calibration technique on the VLBA has not been widespread, the reason being that the data reduction was challenging for the fast temporal variations arising from the troposphere. Nevertheless there are a few examples of such analysis: Rioja & Dodson (2011); Rioja et al. (2014); Marti-Vidal et al. (2016). The Korean VLBI Network (KVN) has introduced a new engineering development in receiver technology (Han et al. 2013) to address this issue, with the capability for simultaneous multi-frequency observations at four bands. Using KVN observations and multi-frequency based calibration techniques it is possible to phase reference the observations at frequencies up to 130 GHz. In this document we will explore some of the scientific topics and current issues that will benefit from this technical capability.

Simultaneous multi-frequency observing enhances mm-VLBI in two main areas: massively increasing the coherence time, for observing weak sources at high frequencies, and enabling accurate astrometric registration between frequency bands. The raw increase in coherence time, even at 130-GHz, has been shown to be easily enlarged from tens of seconds to a few tens of minutes, using observations of the target source only (Rioja et al. 2015). This increase in coherence time would be the equivalent to a three-fold increase in the dish diameter, or an increase of two orders of magnitude in the recorded bandwidth. Furthermore the latter would only apply for continuum science, as this does not assist the important spectral line science cases. The coherence time can be further increased up to several hours by including the observations of a calibrator source. Accurate astrometry is required to allow bona-fide spectral index or Faraday rotation measure investigations in continuum sources and the measurements of the frequency-dependent position of the continuum cores in extragalactic radio sources. Spectral line modes enable high-precision measurements of the physical and chemical conditions of the molecular gas around masing sources, using different tracers.

VLBI is the most developed area driving the demand for simultaneous mm-wavelength receivers, but it is not the only field. Time domain radio astronomy is a growth area in astronomy, and for rapidly changing sources with strong mm-wavelength emission simultaneous mm-wavelength receivers add an important new capability to aid the interpretation of observations. Pulsars are a well-known example and are a target of the BlackHole Cam ERA project, but the example we review in this paper are results from the spectral evolution of the strong and rapidly changing mm emission from the jets in compact X-ray binaries.

Therefore there is a wealth of possibilities opened up by the application of simultaneous multi-frequency mm-VLBI, using demonstrated methods and technology. In this paper we present some of the headline cases for continuum and spectral line VLBI and time-domain observations.

2. Continuum Studies

2.1. Weak Sources

The high angular resolution provided, combined with the opacity effect in high electron density regions that smother the radiation at cm-frequencies, require us to use mm-VLBI to explore the details of the innermost regions of galaxies and AGNs by reaching a spatial resolution of a few tens of Schwarzschild radii (R_s). The Galactic Center Sgr A*, for example, has a R_s of 10 μ -as (Shen et al. 2005), which can only be seen by overcoming the synchrotron self-absorption barriers. Although the physical mechanisms of energetic processes, such as jet formation and acceleration in AGNs have been widely studied at cm-wavelengths, our understanding at mm-wavelengths is still limited due to several considerations, such as the number of available sources, sensitivity, suitable telescopes etc.. Since radio sources become generally weaker as the observing frequency increases, most radio sources are difficult to

detect and image using mm-VLBI, as compared to cm-wavelengths. Most VLBI surveys of continuum sources have therefore been conducted at centimeter wavelengths. In particular, the advent of the VLBA greatly facilitated large VLBI surveys addressing various statistical properties of AGNs; for example, their structure, compactness, brightness temperatures and spectral indices, gamma-ray connection, polarimetry and cosmological evolution (see Table 1). However, weak sources can always be imaged as long as a calibration scheme to compensate for the atmospheric disturbances is included in the experiment. This compensation does not need to be as perfect as for phase referencing, where all errors are corrected, for the simple detection of weak sources. For this, we only require an increase in the coherence time, which, combined with the large bandwidths of modern recording systems, is sufficient for this science case. This technique is referred to as Frequency Phase Transfer (FPT Rioja & Dodson 2011).

As shown in Table 1, there is also a remarkable difference in the number of calibrator sources between cm- and mm-wavelengths. Large calibrator surveys generally aim to extend the database of VLBI sources by finding new sources suitable for use as phase calibrators for astrometry. In the most recent update on the radio fundamental catalog (RFC 2016c) by Petrov (2016), a total of 11,426 compact radio sources has been compiled from S/C/X/K-band observations, and of those more than 3,400 sources are listed in the VLBA calibrator surveys (e.g. VCS1 to VCS6, Petrov et al. 2008 and references therein). In comparison only 121 sources are cataloged at 3 mm (Lee et al. 2008), and these are limited to the very brightest radio sources.

Together with atmospheric fluctuations, the small number of suitable telescopes and their sensitivity at mm-wavelengths make it very challenging to produce high-dynamic range and high-fidelity images of radio sources. If we assume a typical stability of the atmospheric delay in the troposphere as $\sim 10^{-13}$, the expected coherence time for 3 mm VLBI is only about 20 seconds, and the sensitivity of radio telescopes in GMVA stations generally ranges from 1000 to 5000 Jy in system equivalent flux density (SEFD)¹, which leads to a ~ 0.1 Jy 5σ sensitivity in that time. In such situations, VLBI observations using the simultaneous mm-wavelength receivers have demonstrated a novel approach to compensate for phase fluctuations effectively, resulting in an order of magnitude sensitivity enhancement (Jung et al. 2011; Rioja et al. 2015). As a result, a large number of radio sources that have never previously been detected or imaged at mm-wavelengths can now be targeted. This is crucial because our current understanding of physical processes at the centers of galaxies and AGNs has only been studied in a small number of examples, mostly bright radio sources, which may introduce statistical biases.

According to the results achieved by the KVN (where the baselines are short and the atmosphere seems particularly stable), 30 minutes integration of visibilities is possible even at 129 GHz, giving a 5 sigma detection sensitivity as low as \sim 25 mJy. This is shown in Fig. 1 where the detected source SNR is plotted as a function of integration time for no calibration, with FPT calibration, and compared to the theoretical SNR assuming no coherence losses.

Recently, KVN has commenced a Multi-frequency AGN Survey with the KVN (MASK), which is under way in order to densify the grid of available radio sources at 22, 43, 86, and 129 GHz. The early detection statistics for the first 10% of the total sample (~ 1500) with

¹ http://www3.mpifr-bonn.mpg.de/div/vlbi/globalmm/



Fig. 1. The FRING SNR for 3C279 at 129GHz, over a range of integration times, from the single KVN baseline between Ulsan and Tamna, observed on 2012 Apr. 09. Plotted in blue are the direct detections, in red are the detections once calibrated with the scaled phase solutions from 86GHz and in green are the theoretical sensitivities, extrapolated from the SNR achieved in 0.5 min. The FPT calibrated data achieves 77% of the theoretical sensitivity even with an integration time of 30 minutes (Jung et al. 2012).

Survey ID	Wavelength	No. Sources	Reference
CJF survey	$6 \mathrm{cm}$	293	Taylor et al. (1996)
VSOP VLBApls	$6~{ m cm}$	374	Fomalont et al. (2000)
CJF Polarimetry survey	$6 \mathrm{~cm}$	177	Pollack et al. (2003)
ICRF	$3.6~\mathrm{cm}$	~ 500	Ojha et al. (2004, 2005)
			and references therein
MOJAVE	$2~{ m cm}$	>133	Lister & Homan (2005)
2cm Survey	$2 \mathrm{~cm}$	250	Kovalev et al. (2005)
VLBA Calibrator Survey	$13 \& 3.6 {\rm \ cm}$	>3400	Kovalev et al. (2007)
VIPS	$6 \mathrm{~cm}$	1127	Helmboldt et al. (2007)
VERA FSS / $GaPS$	$1.35~\mathrm{cm}$	500	Petrov et al. (2007)
VSOP Survey	$6~{ m cm}$	~ 300	Dodson et al. (2008)
TANAMI	$3.5 \& 1.3 { m cm}$	80	Ojha et al. (2010)
mJIVE-20	$20~{\rm cm}$	>4300	Deller & Middelberg (2014)
GMVA 3mm	$3 \mathrm{mm}$	123	Lee et al. (2008)
ICRF 22 & 43 GHz $$	$13.7~\&~7~\mathrm{mm}$	~100	Lanyi et al. (2010)
KVN Q-CAL survey	$7~\mathrm{mm}$	638	Petrov et al. (2012)

Table 1. Summary of VLBI Surveys, their wavelengths and the number of sources catalogued. The difference in scale of the number of sources in the cm and mm surveys are clear.

a 1 Gbps recording rate (64 MHz bandwidth per band) and using FPT from 22 GHz shows around a 97, 97, 90, and 60% detection rate at 22, 43, 86, and 129 GHz, respectively. This result is very promising for finding many more weak sources by utilizing simultaneous mmwavelength receivers. In addition, it should be noted that simultaneous flux measurements at different frequencies give better constraints on spectral analysis, especially for AGNs that are highly variable on time scales similar to the observation length, and which and whose flares are associated with structural changes in the early stages of their evolution, such as Sgr A* (e.g. Fish et al. 2011).

Nevertheless, it is very difficult to obtain high-fidelity images of compact radio sources, such as the immediate vicinity of supermassive black holes, with the few KVN baselines (max ~500 km), and thus, studying the structure and evolution of (sub-)pc scale emissions is demanding. In addition, the statistical study of relativistic jets using the observed distribution of brightness temperatures (T_b) and/or the intrinsic T_b of mm-VLBI cores will be challenging. Accordingly, increasing the number of mm-VLBI telescopes with simultaneous mm-wavelength receivers is a crucial requirement to deepen our understanding of galaxies and AGNs. With such a capability, mm-VLBI could observe a similar number of sources and obtain comparable sensitivity to that achieved currently with cm-VLBI.

2.2. Astrometry

Astrometry provides fundamental information to a large number of fields in astronomy. An accurate (i.e. astrometric) registration of images (be they total intensity, polarised emission, spectral line or others) obtained at different frequencies is crucial to form a meaningful interpretation based on multi-frequency comparisons, in a similar manner as is required between epochs for multi-epoch temporal studies. Bona fide astrometry can be achieved with a suitable calibration strategy that removes the dominant contribution of the medium through which the signal propagates, while retaining the intrinsic astrometric signature of the source in the phase observable. If so, the astrometric accuracy is ultimately limited by the uncertainty in the precise phase observable and reaches the thermal limit of the instrument ($\sigma_{\rm pos} \sim \theta_{\rm B}$ /SNR), which leads to μ -as astrometry. Phase referencing does this, by interleaving observations of a second source, but this fails for frequencies beyond \sim 43 GHz, defeated by the fast tropospheric fluctuations.

There are a number of image-based, non bona-fide, methods in AGN studies which have been used to align images between frequencies. The core of the jet, defined as the upstream end of the jet, would seem a priori to be the best choice, since it is usually the strongest jet feature. However, opacity effects at centimeter wavelengths can move its position with observing frequency (e.g., Blandford & Königl 1979); an effect known as opacity core-shifts. Furthermore, the blending of multiple components close to the VLBI core at longer wavelengths further complicates the alignment of the images. It is therefore more convenient to choose jet components that are known to be optically thin, strong, and compact, for a more accurate determination of their position. Finding such components is not always easy, specially when comparing observations at relatively distant observing wavelengths, due to the vastly different convolving beams. An alternative method for the alignment of the images can be obtained by performing a cross-correlation of the optically thin jet emission (Walker et al. 2000), which has provided a very simple approach to derive results (e.g., Croke & Gabuzda 2008; Fromm et al. 2013; Molina et al. 2014). However these image-based non-bona fide methods are prone to producing results with reduced accuracy and precision. Hovatta et al. (2014) argued that any results derived without accurate phase referencing are questionable close to the core, which are the regions of most interest in mm-VLBI.

The Korean VLBI Network (KVN) (Kim et al. 2004; Lee et al. 2014) is the first dedicated mm-VLBI array and addresses one of the fundamental limitations of the field, the atmospheric stability, with an innovative multi-channel receiver design. KVN currently consists of three

antennas spread across South Korea, located in the campus of the Universities of Yonsei and Ulsan on the mainland, and in Tamna, on Jeju island. The observing frequencies are centred at 22, 43, 87 and 130 GHz. The multi-channel receiver (Han et al. 2013) of KVN is designed for atmospheric compensation using simultaneous observations at multiple bands, which results in an effective increase of the coherence time, well beyond that imposed by atmospheric propagation. The KVN backend (Oh et al. 2011), combined with the Source Frequency Phase Referencing (SFPR) calibration strategy (Dodson & Rioja 2009; Rioja & Dodson 2011; Rioja et al. 2011, 2014), allows high precision astrometric measurements even at the highest frequencies of 130 GHz (Rioja et al. 2015). We know of no demonstrated upper frequency limit and the method would be expected to work as long as the tropospheric propagation effects were non-dispersive. The baseline lengths between the KVN antennas range between 300 and 500 km, providing a spatial resolution \sim 1 mas at the highest frequency band. Work is on-going to develop a global network of KVN-compatible antennas (Jung et al. 2015) which will provide tens of μ -as resolution.

In Rioja et al. (2014) we presented results of simultaneous SFPR astrometric measurements with KVN at 22 and 43 GHz for continuum sources, along with a detailed comparative study with contemporaneous observations using fast frequency switched SFPR observations, made with the VLBA. Dodson et al. (2014) presented the demonstration for spectral line observations, with the registration of maser emission of H_2O and SiO masers of the AGB star R-LMi at 22 and 43 GHz. In Rioja et al. (2015) astrometric VLBI measurements were successfully extended to all the four frequency bands supported by the KVN, that is up to 130 GHz. Fig. 2 shows the results from this study. More recently Cho et al (in prep) have successfully applied SFPR for registration of the four bands in spectral line observations, comprising the water masers at 22 GHz, and 4 SiO maser transitions at 42.8, 43.1, 86.2 and 129.3 GHz, in VY CMa. These are the first and only cases of astrometry at 130 GHz with VLBI to date.

SFPR is primarily designed for astrometric registration between frequencies, but can produce positional astrometry with respect to an external reference point at high frequencies when combined with conventional phase referencing observations at the lower frequency. An example of such an analysis was the astrometric measurement of R-LMi (Dodson et al. 2014), see Fig. 3. In this case the two decades time span between our observations and the Hipparcos measurements (van Leeuwen 2007) lead to large positional uncertainties. The conventional phase referencing astrometry of the H₂O masers with respect to a quasar, then the SFPR referencing of the emission from SiO masers with respect to that from H₂O masers, allowed the positional astrometry of the star (at 43 GHz) to be determined with significantly better accuracy, reducing the uncertainty in the proper motion for R-LMi by an order of magnitude.

2.3. Faraday Rotation studies

VLBI polarimetric observations are a unique tool to obtain information about the strength, degree of ordering, and orientation of the magnetic field on the plane of the sky in AGN jets. Simultaneous polarimetric VLBI observations at two or more frequencies allow also for the determination of the Faraday rotation of the plane of polarisation, since it is proportional to



Fig. 2. The derived individual core-shifts for a subset of five sources from the S5 polar cap sample, between 22, 43, 86 and 130-GHz. (a)–(e) Polar plots of the decomposed absolute single-source position shifts between two frequencies, for five frequency pairs for the AGNs. These are shown in different colors (K \rightarrow Q (red), K \rightarrow W (blue), K \rightarrow D (black), Q \rightarrow W (green), and Q \rightarrow D (cyan)). In the polar plots the position angles are shown outside the largest circle and are 0° and 90° toward north and east, respectively, and the magnitude units, as specified in the concentric circles, are in μ as. f) shows the expected jet directions. These results have been derived from the SFPR pairwise measurements, using SVD plus the alignment constraint between the jet and the position shift directions, to break the degeneracy. See Fig. 10 of Rioja et al. (2015) for details.



Fig. 3. The astrometric position of the AGB star R-Leo Minoris (small cross), as derived from the centroid of the SiO $J=0\rightarrow 1$ maser emission, with blue (v=1) and green (v=2) contours. The SiO masers are astrometrically registered to the H₂O maser emission (red contours), which is conventionally phase referenced to a known extra-galactic calibrator. The large cross indicates the Hipparcos position for this source, with the bar length being the proper motion error over the 21 years since the optical positional epoch. For details see Dodson et al. (Fig. 5 2014).

the square of the observing wavelength. This can be used to determine the direction of the line-of-sight magnetic field – and therefore the three-dimensional structure of the field, given that the amount of Faraday rotation depends on the line-of-sight magnetic field strength. It provides also a way to probe the thermal plasma around the radio-emitting jet regions, since the amount of rotation depends also on the thermal electron density. This has been used to study, for example, the interaction of the jet with the external medium (Gabuzda et al. 2001; Gómez et al. 2000, 2008).

These Faraday rotation studies are predicated on a proper alignment of the VLBI images across different observing wavelengths, but self-calibration of the visibility data during the imaging process results in the loss of absolute positioning of the source. As discussed previously, registration of the VLBI images is commonly performed by looking for particular optically thin jet features (components) that can be unambiguously identified in all of the images, or be obtained by performing a cross-correlation of the optically thin jet emission. The concerns raised in Hovatta et al. (2014) about the accuracy of these methods, particularly close to the core region, are discussed in Section 2.2. A *bona-fide* astrometric registration of the images can only be obtained through phase-referenced observations, and for mm-VLBI the best method available is SFPR. Nevertheless a number of the results discussed here are based on these non bona-fide astrometry techniques, demonstrating the science that can be extracted.



Fig. 4. Left: RadioAstron images of BL Lac at 1.3 cm from November 11, 2013. Total intensity is shown in contours, linearly polarised intensity in color scale, and white bars indicate the EVPAs. The inset panel shows the uv-coverage. *Right* shows the rotation measure map and a) two-dimensional histograms of the rotation measure and b) Faraday-corrected EVPA in the core area. Adapted from figures in Gómez et al. (2016).

page 10 of 28

Faraday rotation analysis can be used to look for helical magnetic fields in AGN jets, which are expected to arise from the differential rotation of the accretion disk, and are thought to have an important role in the actual jet formation and collimation processes (e.g., McKinney et al. 2013; Zamaninasab et al. 2014). Faraday rotation measure (RM) gradients across the jet width should appear naturally if the jets are threaded by a helical magnetic field due to the systematic change in the net line-of-sight component of the magnetic field across the jet, with increasing values toward the jet boundaries (Laing 1981). The first observational evidence for the existence of such gradients in RM across the jet was given by Asada et al. (2002) based on near-simultaneous multi-wavelength VLBA observations of the quasar 3C 273. Further observations have confirmed the transverse RM gradient in 3C 273 (Zavala & Taylor 2005; Asada et al. 2008a; Hovatta et al. 2012), as well as in other sources (Gabuzda et al. 2004; Asada et al. 2008b, 2010; O'Sullivan & Gabuzda 2009b).

One of the most comprehensives studies of Faraday rotation in AGN jets was performed by Hovatta et al. (2012), in which 149 sources from the MOJAVE sample were observed with the VLBA at four frequencies between 8 and 15 GHz. Four sources, CTA 102, 4C 39.25, 3C 454.3, and 3C 273 were found to contain clear RM gradients across the jet width. The jet in 3C 273 also displayed variations in the RM screen on a time scale of months, which may require some internal Faraday rotation, produced in the jet emitting region.

Studying the magnetic field configuration in the vicinity of the central black holes in AGN jets requires the highest angular resolution possible, which involves VLBI observations at either progressively shorter wavelengths or larger baseline distances, such as those provided by the RadioAstron space VLBI mission. The first polarimetric RadioAstron 1.3 cm observations were performed in November 11, 2013, in which BL Lac was observed in combination with a ground array of 15 antennas (see Fig. 4). Correlated visibilities between the ground antennas and the space radio telescope have been found extending up to a projected baseline distance of 7.9 Earth diameters, yielding a maximum angular resolution of 21 μ as, the highest achieved to date in astronomy (Gómez et al. 2016). Fig. 4 shows the RM image obtained by combining the RadioAstron 1.3 cm observations of BL Lac with simultaneous ground array 7 mm and 2 cm observations. The RM and Faraday-corrected EVPAs in the core area display a clear point symmetry around its centroid. This is better observed in the probability distribution function of the two-dimensional histogram for the RM and Faraday-corrected EVPAs as a function of position angle with respect to the centroid of the core, as shown in Fig. 4. This suggests that the core region in BL Lac is threaded by a large-scale helical magnetic field. as shown in relativistic magnetohydrodynamic simulations (Porth et al. 2011). Whilst we are confident in the solidity of these results, the lack of bona-fide astrometry prevents a robust characterization of the errors associated with the alignment of the images. Furthermore, for other weaker sources a proper astrometric analysis may be required to prevent unambiguous interpretation of the results

There has been significant progress in developing VLBI arrays at progressively shorter wavelengths. The GMVA is now able to provide good quality images in the 3 mm band, with an angular resolution of typically 50–70 $\mu {\rm arcs}$ (e.g., Molina et al. 2014). mm-VLBI offers also the possibility to probe the innermost jet regions, which often appear self-absorbed at centimeter wavelengths. However such multi-wavelength investigations requires accurate registration between wave-bands.

Progress is being made in further improving the resolution through VLBI observations at 1.3 mm making use of the EHT, which includes observations with the PV, PB, LMT, SMTO, SMA, JCMT, and APEX telescopes, and for the first time, the participation of ALMA as a phased array in spring 2017. To honour its name, the EHT is intended to achieve the necessary angular resolution (\sim 10 μ as) to actually discern the silhouettes of the black holes in M87 and SgrA* (Doeleman et al. 2008, 2012; Johnson et al. 2015), and to probe the jet formation regions in other nearby AGN with supermassive black holes (of the order of 10^{9-10} solar masses). Sensitivity is the crucial issue in such investigations, and the increased coherence time provided by SFPR and related methods are highly efficient solutions to this challenge.

Comparison between quasi-simultaneous EHT, GMVA, and RadioAstron observations combining 3 wavelengths (1.3 mm, 3 mm, and 1.3 cm) with almost matching beams will allow us to perform Faraday rotation and opacity analysis with angular resolutions of the order of 20-40 μ as, a ~10-fold improvement with respect to previous studies. However we note that the registration between these different instruments and frequencies will be vital for any robust analysis.

2.4. Opacity core-shifts, γ -ray flares, and the nature of the VLBI core

Results of over seven years of monthly monitoring of 36 blazars (the most luminous and variable BL Lac objects and flat-spectrum radio quasars), with the VLBA at 7mm by the VLBA-BU-BLAZAR program, show that most γ -ray flares are simultaneous within errors with the appearance of a new superluminal component or a major outburst in the core of the jet, which is defined as the bright, compact feature in the upstream end of the jet (Marscher et al. 2008; Jorstad et al. 2013; Casadio et al. 2015). A burst in particle and magnetic energy density is therefore required when jet disturbances cross the radio core in order to produce γ -ray flares, which can naturally be explained by identifying the radio core with a recollimation shock (Gómez et al. 1995; Gómez et al. 1997b; Daly & Marscher 1988).

Multi-wavelength observations of blazars therefore suggest that the radio core is a physical feature (recollimation shock) in the jet at a fixed location. On the other hand, the standard Blandford & Königl conical jet model hypothesises that the core is not a physical feature in the jet, but corresponds to the location at which the jet becomes optically thin, and therefore its position shifts with observing frequency (Blandford & Königl 1979; Königl 1981; Gómez et al. 1993; Lobanov 1998). Multi-frequency VLBI observations at centimeter wavelengths have measured this core frequency shift in multiple sources, albeit without phase-referencing (e.g., Kovalev et al. 2008; O'Sullivan & Gabuzda 2009a; Sokolovsky et al. 2011; Fromm et al. 2015). Furthermore, phase-referenced VLBI observations have shown that in the jet of M87 the radio core indeed corresponds to the optically thick-thin transition between the observing frequencies of 2.3 and 43.2 GHz (Hada et al. 2011).

We have therefore two sets of results, one suggesting that the radio core corresponds to a recollimation shock while the other implies that it marks the transition between the optically thick-thin jet regimes. A possible solution to reconcile these apparently contradicting observational results is to consider that the Blandford & Königl core-shift at centimeter wavelength is produced because the foot of the jet becomes optically thick at these low



Fig. 5. a) A sequence of simulated synchrotron total intensity images computed at 86, 43, 22, 15, 12, 8, and 5 GHz, respectively, using a relativistic hydrodynamical model of a jet with a recollimation shock. b) Position of the simulated core as a function of frequency. The red curve indicates the best fit to the core positions between 5 and 22 GHz, which follows the conical Blandford & Königl jet model. The 43 and 86 GHz data clearly deviate from the opacity core-shift curve, revealing the recollimation shock. c) Astrometric core-shifts of BL-Lac between 4.8 and 43 GHz, plotted as a function of frequency, adapted from Dodson et al. (2016). Overlaid in red is the model from the fitting of the cm-wavelength data (where κ is -0.99 and r_0 is 5.3 mas GHz^{κ})

frequencies. On the other hand, at millimeter wavelength one can see closer to the Black Hole, revealing a recollimation shock that was previously hidden. While this is certainly not the case for M87 (Hada et al. 2011), simultaneous observations at γ /X-rays, optical, infrared, and radio, together with mm-VLBI imaging, have shown that in several radio galaxies and blazars the core indeed is inferred to lie parsecs away from the central black hole (Marscher et al. 2002, 2010; Chatterjee et al. 2011; Fromm et al. 2015).

To test this scenario we have performed numerical simulations using the finite-volume code RATPENAT, which solves the equations of relativistic hydrodynamics (Perucho et al. 2010, and references therein). The jet is launched with a Lorentz factor of 7 and an initial over-pressure of 1.5 times that of the external medium, in order to obtain a recollimation shock. Using the hydrodynamical results as input, we have then computed the synchrotron emission at different observing frequencies (for details of the numerical model used see Gómez et al. 1995; Gómez et al. 1997a; Mimica et al. 2009), adjusting the model parameters so that the jet is optically thin at 86 and 43 GHz, and becomes optically thick at lower frequencies. A magnetic field in equipartition with the particle energy density is assumed. The results are shown in Fig. 5a and b. At cm-wavelengths (5 to 22 GHz) numerical simulations reproduce the opacity core-shift of a Blandford & Königl conical jet model, while at millimeter wavebands (43 and 86GHz) the core position clearly departs from this behaviour, revealing the core as an optically thin recollimation shock at a fixed jet location. Testing this scenario requires astrometric observations at millimeter wavelengths, for which the SFPR technique can provide the necessary high precision (Rioja et al. 2015). Recent observations, see Fig. 5c, suggest that this effect has been detected for the first time, in BL-Lac, reproducing the simulations extremely closely (Dodson et al. 2016).

3. Spectral Studies

3.1. Multifrequency VLBI observations of maser emission in evolved stars

3.1.1. AGB and post-AGB evolutionary phases

Most stars in the sky, with initial masses between about 0.5 and 8 M_{\odot} , reach the AGB phase (or a similar one) at the end of their lives. After this phase, most of the initial mass has

been ejected to form planetary nebulae (PNe), after the short phase of protoPN. Finally, this ejected material will return to the interstellar medium, enriching it (and the new generations of stars to be formed from it) with heavy elements.

The copious mass-loss of AGB stars forms thick circumstellar envelopes (CSEs) around them. Mass-loss is a basic phenomenon in their evolution, the ejection rate tends to increase with time and, by the end of the AGB phase, it is so strong, close to $10^{-3} M_{\odot} \text{ yr}^{-1}$, that most of the initial stellar mass is ejected in a relatively short time. Then, the AGB phase ends, the stellar core becomes exposed and can be seen, becoming the new central star. This new star is very compact and shows an increasingly high temperature, rapidly evolving to the blue and white dwarf phase.

The circumstellar nebula is also evolving very fast: from the nearly spherical CSE in the AGB phase, which is in relatively slow expansion (at, say, 10-15 km s⁻¹), to the strongly axisymmetric PNe around the dwarfs, which show very fast bipolar outflows, with axial velocities as high as several hundred km s⁻¹, (Bujarrabal et al. 2001; Balick & Frank 2002, etc). This metamorphosis is a very fast and spectacular phenomenon. Within about 1000 yr, the star becomes a blue dwarf able to significantly ionize the nebula, which has already developed a wide bipolar shape.

The mass-loss process in AGB stars behaves, basically, in two phases, e.g. Höfner et al. (2003). In inner circumstellar regions, the stellar pulsation (all these old stars are strongly pulsating, in more or less regular modes) propagates in the outer atmospheric layers, leading to a smooth distribution of the atmosphere and keeping quite high densities at relatively long distances from the photosphere. When the gas temperature drops to about 1000 K, at a few photospheric radii, the formation of dust grains becomes very efficient and refractory material condenses. Radiation pressure acts very efficiently on dust grains, which are dynamically coupled with gas. The result is that the whole circumstellar layers start expanding, quickly reaching (at 10-20 stellar radii) their final expansion velocity. The circumstellar dynamics, which is in fact driving the evolution of the AGB stars and the ejection of PNe, is really active only in those inner regions. The study of the inner layers of CSEs is therefore crucial to understand these phases of the stellar evolution.

3.1.2. SiO and H_2O masers from AGB stars

SiO (at 43, 86, 130, 215, ... GHz) and H_2O (22 GHz) masers are known to be very useful tools for studying the detailed structure and dynamics of the AGB CSEs. SiO masers come from regions at about 2-4 stellar radii, where pulsation is propagating and dust is not yet completely formed. The SiO maser lines in general come from a ring of spots, centered on the star, very probably due to the dominant tangential amplification (Diamond et al. 1994; Desmurs et al. 2000, etc). On the other hand, H_2O masers form shortly after grain formation, when the final expansion velocity is being reached, e.g. Richards et al. (2012). The very high angular resolution of VLBI imaging at these frequencies allows very detailed mapping of such crucial shells, providing also valuable information on their dynamics. However, the complex pumping of the masers, which is not well understood (particularly for SiO), and the often poor information about the relative positions of the spots detected at the different frequencies strongly limit our studies of circumstellar masers from VLBI data.



Fig. 6. Observations of SiO J=1-0 v=1 (greys) and v=2 (contours) in IRC+10011, (Fig. 9 Soria-Ruiz et al. 2004). Note that both lines occupy similar regions but that, in fact, the spots are never coincident, in spite of the uncertain relative astrometry.

SiO masers appear in rotational transitions (J=1-0, J=2-1, J=3-2, ...) within vibrationally excited states (v = 1, 2, 3, ...). The various proposed pumping models agree in identifying the inversion mechanism: line trapping in the rovibrational transitions $(v \rightarrow v-1)$, which decreases the original radiative probabilities to efficiency probabilities that are relatively lower for higher-J levels; see e.g. Bujarrabal & Nguyen-Q-Rieu (1981); Lockett & Elitzur (1992); Gray et al. (2009). This leads to a systematic overpopulation of levels with relatively high J-values and to the inversion of the $J \rightarrow J - 1$ transitions in the v > 0 states, which

is particularly efficient for lower frequency lines. What is not yet well known is the source of the energy, i.e. the way in which we excite molecules from the ground v=0 state, which can be collisional or radiative (Humphreys et al. 2002; Bujarrabal 1994). The issue is important, because the relationship between the physical conditions for the maser emission must depend on the pumping mechanism. If we wish to probe the physical conditions based on the observed radiation we must understand how the former generates the latter. Radiative pumping tends to discriminate more clearly the conditions required for the pumping of the different v-states. The fact that the maser spots of the v=1 and v=2 J=1-0 lines tend to form clusters that are similar for both lines (the ring radius being slightly smaller for the v=2 line) has been argued to favour collisional pumping (Miyoshi et al. 1994; Desmurs et al. 2000; Rioja et al. 2008; Kamohara et al. 2010, etc). The J=1-0 masers are intense and their frequencies, around 43 GHz ($\lambda = 7$ mm), are easily reachable at present in VLBI, therefore, they are the best observed SiO masers. However, high-resolution (<mas) observations have shown that, curiously, the spots of the two lines are practically never coincident, which is interpreted as supporting radiative pumping (Fig. 6). Of course, such a comparison is hampered by the often poor bona-fide astrometry in VLBI experiments at 7mm wavelength. Of those listed above only Rioja et al. (2008) and Kamohara et al. (2010) were phase referenced, and both observed with VERA. At higher frequencies (higher J-values), VLBI phase referencing experiments are more difficult and astrometry is still not yet possible (although we note that a significant number of four-band KVN observations, albeit with greater than mas resolutions, are in preparation, and that the first KaVA results have been recently published. Yun et al. (2016)).

The problem also appears when comparing H_2O with SiO masers. SiO masers are clearly placed in a more central region, but we are not sure exactly how they are placed in relation to the H_2O spots, as these in general show a complex distribution with a less obvious center. When the transitions are astrometrically registered, however, this issue is addressed, as in Dodson et al. (2014).

The publication of maps of other SiO lines has raised new problems. The v=3 J=1-0 line is placed at more than 5000 K over the ground, requiring a high excitation, and one expects it to be placed in a different region, probably closer to the star. But observations place it at about the same distance, in very similar distributions to that for v = 1 and 2 and always with very rare coincidences at the smallest scale. On the other hand, the v=1 J=2-1 maser, which theoretically shares the same pumping mechanism and excitation requirements as v=1 J=1-0, appears in clearly different regions and at further distances than the 7mm lines. These new observations seem in clear contradiction with all the simple pumping models. Needless to say, the v=3 J=1-0 maser is weaker than the v=1,2 J=1-0 ones and the v=1 J=2-1 line frequency, 86 GHz, is hard to observe in VLBI, therefore, astrometry in such observations has hitherto been impossible.

A possible solution to the puzzle is the presence of line overlap. Photons emitted by a rovibrational transition of H₂O, $v_2=1$ 11_{6,6} – v=0 12_{7,5}, can be absorbed by the v=1 J=0 – v=2 J=1 component, because both frequencies are almost exactly the same (Olofsson et al. 1981; Soria-Ruiz et al. 2004). It can be shown that H₂O often emits strongly at that frequency in our sources. Therefore, the absorption of this radiation by SiO leads to an underpopulation of v=1 J=0 and an overpopulation of v=2 J=1 line, quenching the page 16 of 28

(otherwise expected to be intense) v=2 J=2-1, which is known to be anomalously weak; in fact the effects of overlap were first invoked to explain this observational fact. At the same time, this phenomenon introduces a strong over-excitation of the two 7mm masers, v=1,2 J=1-0, whose excitations are now strongly coupled. Calculations show that the excitation conditions of both lines are now very similar and also similar to those of the v=3 J=1-0 line (Imai et al. 2010; Desmurs et al. 2014). Meanwhile, v=2 J=2-1 now appears to require different conditions, in particular lower densities that should appear at larger distances from the star. Radiative models including line overlap can explain, at least qualitatively, all the observational results. However, we stress the scarce available data with high sensitivity and with any meaningful astrometric registration, particularly at high frequencies. We do not yet have accurate and systematic information on the relative distributions of these lines, for which not only do we need good absolute astrometry but we would also need to recover (almost) all the maser flux, to better understand the complete brightness distribution. In particular, we lack VLBI data of lines at higher frequencies, necessary to be sure that line overlap is the main (or unique) major modification of the basic pumping mechanisms.

3.1.3. Maser lines in proto-PNe

As soon as the star+nebula system leaves the AGB phase, the observation of the masers associated with them becomes more and more difficult. The reason is that the inner circumstellar masers are becoming more and more diffuse at the same time as the mass-loss rate decreases. SiO masers are very rarely observed. There is however a paradigmatic example showing both SiO and H₂O masers: the strongly bipolar nebula OH 231.8+4.2, the Calabash Nebula.



Fig. 7. Observations of SiO and H_2O masers in OH 231.8+4.2, compared with the optical image. The uncertainty in the SiO maser position is about 200 mas, too large to definitively determine its relationship with the H_2O clumps. From Desmurs et al. (Fig. 1 2007)

We have seen the spectacular evolution from the AGB to the PN phase. It is thought that the strong axial symmetry typical of post-AGB nebulae is due to the ejection, during the first protoPN phases, of very fast and collimated stellar jets. These jets shock the fossil slow-moving AGB envelope, generating a series of axial shocks that cross the massive CSE, inducing high axial velocities in it (see e.g., Balick & Frank 2002; Bujarrabal et al. 2001). The very large amounts of energy and momentum observed in young PNe impose severe limitations on the jet launching mechanism. At present, the only way to explain the origin of such energetic flows is to assume that a fraction of the ejected CSE is reaccreted by the central star or a companion through a rapidly rotating disk (Soker 2001; Frank & Blackman 2004). The jet would then be powered by a magnetocentrifugal launching process, similar to that at work in forming stars. For that to be efficient, the presence of a stellar companion (or at least a massive planet) is necessary, since otherwise the circumstellar material lacks the angular momentum to form an accretion disk. OH 231.8+4.2 is strongly bipolar, shows a very fast and massive bipolar outflow, and is known to harbor a binary star in its center. It is therefore a textbook candidate to study these crucial phenomena.

VLBI observations of SiO and H₂O masers yielded very promising results (see Sánchez Contreras et al. (2002), Desmurs et al. (2007), and Fig. 7). Water vapor emission comes from two regions in opposite directions along the nebula axis and their velocity field is fully compatible with the general velocity field in that direction, suggesting that the H_2O clumps represent the inner nebula, at the base of the bipolar flow. SiO masers occupy smaller regions and are placed almost exactly perpendicular to the axis. The movements depicted by the observations of SiO are compatible with a disk orbiting the central star(s). In principle, we are seeing in this object the whole central structure of disk plus outflow that we would like to see to confirm our ideas on the post-AGB nebular dynamics, which has not been observed to date in other sources. But the relative astrometry is poor and we are not sure that really the SiO structure is placed at the center of the nebula, between the two H_2O clusters. The problem is then serious, as we lack this crucial datum to draw definitive conclusions on the disk/outflow association. For the last 10 years we have been trying to improve the astrometry, but, up to now, the best SiO maser position obtained is uncertain by about 200 mas, completely insufficient for our purposes. Furthermore the SiO masers in this source have been fading over the past several years, so new VLBI measurements are more and more difficult. The VLBI data on OH 231.8+4.2 constitute a self-explanatory example of the effects of the lack of accurate and sensitive VLBI data in our studies of the inner post-AGB nebulae.

3.1.4. Conclusions

We have seen how important the VLBI observations of various maser lines are in the study of the most interesting layers of nebulae around AGB stars: the inner regions in which the dynamics are still active and the relevant phenomena to understand these evolutionary phases are actually taking place. In AGB stars, they harbor the grain formation and gas+dust acceleration layers; in post-AGB objects, this inner nebula includes the outflow acceleration regions and may hold the key to understanding the formation and shaping of PNe. In both cases, however, the existing VLBI data are still insufficient to address these problems, because of the lack of systematic observations with accurate astrometry and high sensitivity. Good VLBI data are also necessary to understand the SiO maser mechanism, which is still under debate.

Accurate astrometry is still difficult at 7mm. At higher frequencies, VLBI experiments are very difficult and conventional astrometry is impossible. In all cases, sensitivity limitations are significant and an important fraction of the total flux is lost in the interferometric process. Thanks to phase transfer between the SiO lines and from 22 GHz, sensitive multifrequency observations will strongly help to map several transitions, out to the sub-mm regime. They will also help to systematically obtain good relative astrometry for the different transitions.

3.2. Massive Stars

3.2.1. Introduction

The formation of isolated low-mass (M $\sim 1~M_{\odot}$) stars is now understood quite well. It proceeds through (1) mass accretion onto the protostar through a Keplerian disk, an expected result of angular momentum conservation, and (2) ejection of material via a jet collimated along the disk axis, which removes angular momentum from the disk allowing matter to accrete onto the star. Jets are generally modeled as magneto-centrifugally driven winds, powered by the rotation and gravitational energy and channeled along the magnetic field lines either from the disk inner edge ("X-wind", Shu et al. 1995), or across a much larger (up to 100 AU) portion of the disk ("Disk-Wind", Pudritz & Banerjee 2005).

The formation of more massive stars might require substantially different accretion/ejection processes. For the mass accretion rates predicted by theory (e.g. Shu 1985), the accretion time becomes longer than the protostar contraction time for masses $\geq 8~M_{\odot}$. At this point, the star reaches the Zero Age Main Sequence (ZAMS) and, as a consequence, its radiation pressure would be sufficient to halt the inflow (Wolfire & Cassinelli 1987), thus preventing further increase of the stellar mass. A number of scenarios have been put forward to solve this "radiation pressure" problem for high-mass star formation: much higher mass accretion rates, competitive accretion (Bonnell et al. 2003) and/or stellar mergers (Bonnell et al. 1998) in clusters.

However, recent 3-D, radiation hydrodynamic simulations of high-mass SF seem to indicate that accretion disks and collimated outflows could also be an effective mechanism for assembling very massive stars, up to \approx 140 M_{\odot} (e.g. Kuiper et al. 2010). The disk geometry focuses the accretion flow onto the equatorial plane, allowing the accreting material to overcome the strong radiation pressure and reach the stellar surface. These models also predict beaming of the stellar photons into the lower-density outflow cavity, which helps to alleviate the radiation pressure in the equatorial plane. The size of accretion disks around high-mass YSOs is predicted to be on the order of only a few 100 AU, whereas clustering of multiple forming stars should occur on scales of 10^3 AU (e.g. Krumholz et al. 2009).

From all the above it is clear that to improve our understanding of the accretion and ejection processes in high-mass YSOs, it is essential to image linear scales as small as 10–100 AU (requiring angular resolutions <0.1 at the typical distances > 1 kpc of massive star forming regions) to resolve the gas kinematics around single objects. To this purpose very useful diagnostic tools are the intense molecular (in particular SiO, H₂O, and CH₃OH) masers often observed in proximity to high-mass YSOs. Thanks to their high brightness temperature ($\geq 10^9$ K), molecular masers can be targets of VLBI observations, which, achieving submilliarcsec angular resolutions, permit us to derive the proper motions of the maser "spots"

(i.e., the single maser emission centers) and access the full 3-D gas kinematics. However for exact reconstruction of the kinematics the different observations have to be astrometrically registered. The only way to register mm-VLBI reliably is with the SFPR method, and the best observing efficiency for SFPR is provided with simultaneous mm-VLBI receiver systems (Rioja et al. 2014). Combining maser VLBI data with (sub-arcsecond) interferometric observations of thermal (continuum and line) emissions, one can get a very detailed view of the gas kinematics and physical conditions near the forming star. Such a study has been performed so far only towards a small number of objects (see discussion in Moscadelli et al. 2011). In the following we illustrate one of the most remarkable results of maser VLBI applied to high-mass SF.



Fig. 8. Maser kinematics in Source I. Left Panel) Top: SiO v=0 and H₂O masers (*open circles* and *squares*, respectively – (Fig. 3 Greenhill et al. 2013)), velocity-integrated SiO v=1 masers (*red contours* – Goddi et al. (2009)), and 7 mm continuum (*black contours* – Reid et al. (2007)), as mapped with the VLA. Middle: expanded view of SiO v=0 maser proper motions in the northeastern lobe of Source I. Bottom: expanded view of the southwest lobe. Colors denote maser $V_{\rm LSR}$ as indicated in the wedge on the bottom left corner of the bottom panel. Right Panel) Combined (velocity-integrated) SiO v=1 and v=2 maser emission distribution (*red-tone image*) observed with the VLBA (Fig. 5 Matthews et al. 2010), and 7 mm continuum (*color map*) imaged with the VLA (Reid et al. 2007).

3.2.2. Source I in Orion BN/KL

The closest (415 pc) and best studied high-mass YSO is Source I in the Orion BN/KL region. The continuum emission of Source I, imaged at 7 mm with the Very Large Array (VLA) by Reid et al. (2007), is highly elongated and consistent with an accretion disk ionised by a $_{page 20 of 28}$

YSO of $\approx 10 \ M_{\odot}$. This source has been monitored by Matthews et al. (2010) with the Very Long Baseline Array (VLBA) in vibrationally-excited SiO maser transitions every month for over three years, and a movie of the 3-D molecular gas flow was created. The SiO masers are distributed symmetrically around the ionised disk, outlining four radial arms connected by two tangential bridges. The pattern of SiO maser proper motions shows that material in the bridges rotates about the disk axis, whereas gas in the arms streams radially away from the star. Thus, the vibrationally-excited SiO masers are tracing both a compact rotating disk and a wide-angle wind emanating from the disk at radii < 100 AU (Fig. 8, right panel), which can be modeled in terms of a magnetocentrifugally driven wind (Vaidya & Goddi 2013). VLA imaging of ground-state SiO maser transitions, probing larger scales, shows that the wide-angle wind collimates into a bipolar outflow at radii of 100-1000 AU (Fig. 8, left panel) (Greenhill et al. 2013). This study has provided direct evidence for the formation of a massive star via disk-mediated accretion and revealed for the first time the launch and collimation region of an outflow from a rotating compact disk on scales comparable with the Solar system. SiO masers are ideal candidates for SFPR astrometric registration, as the frequency ratio between the transitions is extremely close to integer. Integer frequency ratio significantly simplifies SFPR, although with care one can succeed using non-integer ratios (Dodson et al. 2014).

3.2.3. Advantages of multi-frequency maser observations

High-frequency, single-dish studies have shown that the commonly observed, centimeter wavelength, masing molecules emit many masing transitions at millimeter wavelengths, too. For instance, for methanol, 16 distinct maser transitions have been detected over the frequency range 6-240 GHz (see, for instance, Ellingsen et al. 2012), and for water, the 14 (so far observed) maser lines cover the frequency range 22-660 GHz (see, for instance, Neufeld et al. 2013). These "millimeter wavelength" masers open up new perspectives for star-formation studies as, in general, for a given molecule, models of maser pumping predict that transitions at different frequencies are strongly inverted and become intense masers under varying physical and chemical conditions (described, in particular, in terms of (n_{H_2}) density, gas and dust temperature, molecular abundance; see, for instance, Neufeld & Melnick (1991), for water, and Cragg et al. (2005), for methanol masers). As an example, Fig. 9 shows the dependence of the brightness temperature of four different methanol masers on various physical parameters, as predicted by the models of Cragg et al. (2005). The intensity ratios of different maser transitions from the same molecule can thus be used to probe the physical and chemical conditions of the gas, as long as the observations are sufficiently well registered to ensure that the measurements arise from the same emission regions. The precision with which the physical parameters of the gas can be determined with this method clearly increases with the number of observed maser transitions. To this purpose the role of the newly-detected millimeter wavelength masers will be fundamental, as will the precision alignment of the observations in different wave bands. VLBI observations of maser emissions over an extended frequency range could in principle allow us to map the density and temperature of the gas surrounding the high-mass YSO with milliarcsecond accuracy. Whilst these observations hitherto have not been observed simultaneously, given that we can now register different transitions together using the SFPR method, such capabilities



Fig. 9. Effect of parameter (gas and dust temperature, gas and specific column density) variations on the brightness temperature of four different methanol masers. The four curves are labelled with the frequency of the maser transition to which they refer: 6.6, 12.1, 23.1 and 107 GHz (reproduced from Fig. 2 in Cragg et al. (2005)).

would greatly enhance the scientific return from the observations. The basic assumption of the maser models, which can be observationally verified, is that the various maser lines are effectively emerging from the same volume of gas. If applied to a large enough sample of sources, multi-frequency maser VLBI, by constraining in a consistent way the physical conditions and kinematics of the gas (see below) in the proximity of (within 10–100 AU from) the high-mass YSOs, can be very useful to track the evolutionary phase of the high-mass YSOs, which is presently poorly determined owing to insufficient angular resolution and/or ambiguous interpretation of the observables.

3.2.4. Prospects for the Future

Maser VLBI offers a unique tool to investigate high-mass SF at the smallest (~10 AU) scales, unaccessible even to the new generation of millimeter interferometers. In the few objects studied so far using both maser VLBI and thermal interferometric observations, the maser 3-D kinematics and the $V_{\rm LSR}$ distribution of the (thermal) tracers of high-density gas complement each other, providing a picture of the gas motion around the forming star at radii from 10 to 10^4 AU. The results obtained seem to indicate that disks and jets could play an

important role in the formation of more massive stars, as well. However, maser VLBI reveals also some new interesting kinematic features, whose significance for the stellar formation process has still to be deciphered.

In the case of maser emission, owing to the constraints imposed by the maser pumping, one can expect that a particular region of gas around the YSO offers the most suitable conditions for producing maser action in different lines (from the same molecule). In general, this expectation has been mostly confirmed by previous multi-line VLBI observations, but recent interferometric studies have also demonstrated that maser lines of very different excitation can effectively be used to trace different physical conditions. As an example, Hirota et al. (2014) have recently performed ALMA observations of two millimeter, water lines, the 321 GHz (1862 K above the ground state) and the 336 GHz (at 2956 K) lines, towards Source I in Orion KL. While the spatial and velocity distribution of the 321 GHz emission is elongated and traces the root of the outflow close to the surface of the accretion disk, the 336 GHz line has a compact structure and is most likely emerging from hot (≈3000 K) neutral material orbiting the YSO in an edge-on small (radius of 50 AU) ring. Therefore, VLBI observations of different maser lines from the same YSO could turn to be even more strategic for kinematic studies, if the spatial associations are registered to determine the physical parameters of the gas.

4. Time domain observations of X-ray binary jets

Black hole X-ray binaries (BH XRBs) show much of the same physics as observed in AGN, exhibiting relativistic jets powered by an accretion flow. However, while they are typically factors of 10^2-10^5 times closer than nearby AGN, their black hole masses are 10^5-10^8 times smaller. This implies that the physical size scales that we can probe with high-frequency VLBI correspond to a larger number of gravitational radii in BH XRBs than in AGN. Nonetheless, since timescales close to a black hole scale with mass, BH XRBs afford us the unique advantage of studying the physical processes of jet launching and evolution in real time.

In the current phenomenological picture describing XRB outbursts (Fender et al. 2004), compact, steady jets are seen in the hard X-ray spectral state at the beginning and end of an outburst. Such compact jets have size scales of a few AU, and using VLBI have been directly resolved along the jet axis in two systems (Dhawan et al. 2000; Stirling et al. 2001). While these compact jets are believed to have a relatively low Lorentz factor ($\Gamma < 2$), *this has never been directly measured*. No counterjets have ever been definitively imaged in hard state compact jets, so the jet/counterjet brightness ratio cannot be used to determine the jet Doppler factor.

At the peak of the outburst, there is a transition to a softer X-ray spectral state, upon which discrete, transient ejecta are launched. These are much brighter than the compact jets, and move away from the central source with proper motions of up to tens of milliarcseconds per day, in some cases showing apparent superluminal motion (e.g. Mirabel & Rodríguez 1994; Tingay et al. 1995). While the proper motions of the ejecta can be used to place constraints on fundamental jet parameters such as speed, inclination angle, and the associated Lorentz and Doppler factors (e.g. Mirabel & Rodriguez 1999), the typical large uncertainties in source distance imply that we can only place lower limits on the true Lorentz factors (Fender

2003). VLBI studies of these transient ejecta are complicated by the strong changes in jet flux density and morphology over the course of an observing run, which violate the fundamental assumptions of aperture synthesis. Such issues become a particular problem at mm-wavelengths, where the amplitude variability is stronger and the angular resolution is higher. However, for typical BH XRBs the jet structure is relatively simple, being made up of a few unresolved components. Thus, an observation can be broken up into multiple smaller chunks, which do not individually violate the key tenets of interferometry, and can be imaged and analysed separately (e.g. Dhawan et al. 2000). Alternatively, high time resolution photometry at multiple radio frequencies can be used to glean some information on the jet structure. The characteristic variability timescale at any given frequency provides a measure of the jet size scale at the surface where the optical depth is unity at that frequency (i.e. the radius of the jets). As an example, Miller-Jones et al. (2009) used the minimum variability timescale seen in simultaneous 15 and 43 GHz VLA observations of Cygnus X-3 to determine the characteristic size of the jets during a period of minor flaring, probing scales of 2–4 AU — significantly smaller than that which can be resolved with VLBI.

The *combination* of both a resolved radio jet and sensitive, high time-resolution photometry at multiple radio frequencies, together with simultaneous X-ray observations, can provide an enhanced and unique probe of the structure and geometry of BH XRB jets. As the X-ray variability from the accretion flow propagates downstream in the jet, we can observe it being manifested in the jet. Since we see emission from the surface of optical depth unity at any given frequency, this is seen first at optical/infrared frequencies from closest to the jet base, and moves to lower frequency with time as the fluctuations propagate downstream. This has already been demonstrated via correlations between optical/infrared and X-ray variability (Casella et al. 2010; Kalamkar et al. 2016), but not to date in the mm or radio bands. With simultaneous, high-sensitivity observations at multiple mm wavelengths (where the variations are stronger and less smoothed out than in the cm band), we can determine the time delay between frequencies as a perturbation propagates downstream. If the characteristic height along the jet is also known from the VLBI observations, along with the exact launch time from simultaneous X-ray observations, these then give a direct measure of the jet speed. Together with constraints on the jet radius from the characteristic variability timescales at the different frequencies, we can also probe the jet geometry, in particular the opening angle, which is typically unknown in BH XRBs (Miller-Jones et al. 2006). Thus, simultaneous observations at multiple mm bands can provide one of the few feasible means for directly measuring the jet speed and geometry.

We have started trials of this VLBI+multi-band photometry technique using observations of bright transient ejection events from the 2015 outburst of the BH XRB V404 Cyg (Miller-Jones et al., in prep; Tetarenko A. et al., in prep.). The combination of VLBI imaging and Markov chain Monte Carlo modeling of our multi-frequency light curves (Fig. 10) is providing a powerful probe of the geometry, speed, structure, and energetics of the discrete ejecta that were launched during these flaring events. This technique will subsequently be extended to the compact jets seen in sufficiently bright BH XRB outbursts. However, the existing observations have required a combination of the VLA in sub-array mode to provide the multi-frequency photometry, and the VLBA to provide the imaging. The adoption of a simultaneous multi-wavelength mm-VLBI capability would enable such observations on a single instrument,



Fig. 10. Simultaneous multi-frequency light curves of the 2015 outburst of the BH XRB V404 Cygni, taken using the VLA in sub-array mode. Rapid variability corresponding to the ejection, expansion and fading of multiple jet knots (directly imaged at 2 cm with the VLBA; Miller-Jones et al. in prep.) is observed. The higher-frequency light curves peak earlier, and at higher flux densities, and are less smoothed out than their counterparts at lower frequencies. This trend continues all the way up to the sub-mm, as evidenced by JCMT observations at frequencies as high as 666 GHz (Tetarenko et al. 2015).

thereby removing both the double-jeopardy and logistical complications involved in scheduling strictly-simultaneous observations on two different facilities. Clearly this combination of high-resolution spatial and time domain analysis will benefit immeasurably from observations with continuous, contemporaneous mm-wave systems such as that provided by the KVN, once the capability is extended to longer and more sensitive global baselines.

5. Concluding Remarks

We have presented a few highlights of the science cases which would be advanced by the wide-spread adoption of recording systems capable of allowing simultaneous mm-wavelength VLBI. These are summarised in Table 2.

The science goals cover a range of spectral line and continuum VLBI science, such as AGN polarimetric studies, and alignment of maser transitions to probe different physical conditions around massive star forming regions and AGB stars. Registration between the different frequencies is essential for the scientific interpretation in these fields. New methods and technologies have allowed for registration of the mm-wavelengths. Similarly, time-sensitive observations of multiple frequencies in, for example, X-ray binaries are best delivered by systems capable of observing a wide range of frequencies simultaneously. There are a multi-

tude of possible technological solutions that could deliver these capabilities, and these were explored in the ERATec meeting, October 2015 (Bartolini et al. 2016). VLBA style fast switching has been demonstrated successfully for VLBI, but has significant observational overheads and is very unlikely to succeed at frequencies greater than 86GHz. KVN style recievers are large but have much reduced overheads and have worked at all KVN frequencies (i.e. upto 130GHz). The results from the KVN in all VLBI science fields are being published now, with a special issue of the Journal of Korean Astronomical Society (2015, Vol. 48, No. 5) providing a review. Time domain observations for X-ray binaries have been recently performed. Sub-arraying solutions are being investigated at the Australia Telescope Compact Array. These have demonstrated mm-band phase stabilisation and are expected to successfully demonstrate mm-band frequency phase referencing shortly, once the polarisation issues are resolved.

The cases presented in this paper represent the benefits that will accrue to existing scientific projects. The outcome of new technologies is always that they enable new measurements and open new scientific applications, so those listed here will only be a subset of the expected fields of study.

Science Case (section)	Domain	Requirements		
Weak Sources (2.1)	Continuum VLBI	mm-band phase stabilisation		
Spatial Astrometry (2.2)	VLBI	mm-band phase referencing		
Faraday Rotation Studies (2.3) &	Continuum VLBI	mm-band frequency		
Spectral Index (2.4)		phase referencing		
Alignment of Maser transitions	Line VLBI for Star-forming	mm-band frequency		
(3.1 & 3.2)	& Evolved Star regions	phase referencing		
X-ray Binaries (4)	Temporal studies	Simultaneous mm-band observing		

Table 2. Summary table for the science goals described here

Acknowledgements. Figure 1 is reproduced by permission of POS. Figures 2 to 5 and 8 are reproduced by permission of the AAS. Figure 6 and 7 are reproduced by permission of A&A. Figure 9 is reproduced by permission of MNRAS.

References

- Asada, K., Inoue, M., Kameno, S., & Nagai, H. 2008a, ApJ, 675, 79
- Asada, K., Inoue, M., Nakamura, M., Kameno, S., & Nagai, H. 2008b, ApJ, 682, 798
- Asada, K., Inoue, M., Uchida, Y., et al. 2002, PASJ, 54, L39
- Asada, K., Nakamura, M., Inoue, M., Kameno, S., & Nagai, H. 2010, ApJ, 720, 41
- Asaki, Y., Saito, M., Kawabe, R., Morita, K.-I., & Sasao, T. 1996, Radio Science, 31, 1615
- Balick, B. & Frank, A. 2002, ARAA, 40, 439
- Bartolini, M., Bolli, P., Keller, R., et al. 2016, Colle di Galileo, 5, 33
- Blandford, R. D. & Königl, A. 1979, ApJ, 232, 34
- Bonnell, I. A., Bate, M. R., & Vine, S. G. 2003, MNRAS, 343, 413
- Bonnell, I. A., Bate, M. R., & Zinnecker, H. 1998, MNRAS, 298, 93
- Bujarrabal, V. 1994, A&A, 285
- Bujarrabal, V., Castro-Carrizo, A., Alcolea, J., & Sánchez Contreras, C. 2001, A&A, 377, 868
- Bujarrabal, V. & Nguyen-Q-Rieu. 1981, A&A, 102, 65
- Carilli, C. L. & Holdaway, M. A. 1999, Radio Science, 34, 817
- Casadio, C., Gómez, J. L., Jorstad, S. G., et al. 2015, ApJ, 813, 51
- Casella, P., Maccarone, T. J., O'Brien, K., et al. 2010, MNRAS, 404, L21
- Chatterjee, R., Marscher, A. P., Jorstad, S. G., et al. 2011, ApJ, 734, 43
- Cragg, D. M., Sobolev, A. M., & Godfrey, P. D. 2005, MNRAS, 360, 533
- Croke, S. M. & Gabuzda, D. C. 2008, MNRAS, 386, 619
- Daly, R. A. & Marscher, A. P. 1988, ApJ, 334, 539
- Deller, A. T. & Middelberg, E. 2014, AJ, 147, 14
- Desmurs, J.-F., Alcolea, J., Bujarrabal, V., Sánchez Contreras, C., & Colomer, F. 2007, A&A, 468, 189
- Desmurs, J. F., Bujarrabal, V., Colomer, F., & Alcolea, J. 2000, A&A, 360, 189
- Desmurs, J.-F., Bujarrabal, V., Lindqvist, M., et al. 2014, A&A, 565, A127
- Dhawan, V., Mirabel, I. F., & Rodríguez, L. F. 2000, ApJ, 543, 373
- Diamond, P. J., Kemball, A. J., Junor, W., et al. 1994, ApJL, 430, L61
- Dodson, R., Fomalont, E. B., Wiik, K., et al. 2008, ApJS, 175, 314

page 26 of 28 $\,$

- Dodson, R., Rioja, M., Molina, S., & Gómez, J. 2016, Submitted Dodson, R. & Rioja, M. J. 2009, VLBA Scientific Memorandum n. 31: Astrometric calibration of mm-VLBI using "Source/Frequency Phase Referenced" observations, Tech. rep., NRAO

- Dodson, R., Rioja, M. J., Jung, T.-H., et al. 2014, AJ, 148, 97 Doeleman, S. S., Fish, V. L., Schenck, D. E., et al. 2012, Science, 338, 355 Doeleman, S. S., Weintroub, J., Rogers, A. E. E., et al. 2008, Nature, 455, 78 Ellingsen, S. P., Sobolev, A. M., Cragg, D. M., & Godfrey, P. D. 2012, ApJL, 759, L5 Ender, B. P. 2002, MNRAS, 240, 1352
- Fender, R. P. 2003, MNRAS, 340, 1353
- Fender, R. P., Belloni, T., & Gallo, E. 2004, MNRAS, 355, 1105
- Fish, V. L., Doeleman, S. S., Beaudoin, C., et al. 2011, ApJL, 727, L36 Fomalont, E. B., Frey, S., Paragi, Z., et al. 2000, ApJS, 131, 95 Frank, A. & Blackman, E. G. 2004, ApJ, 614, 737

- Fromm, C. M., Perucho, M., Ros, E., Savolainen, T., & Zensus, J. A. 2015, A&A, 576, A43 Fromm, C. M., Ros, E., Perucho, M., et al. 2013, A&A, 557, 105
- Gabuzda, D. C., Murray, É., & Cronin, P. 2004, MNRAS, 351, L89
- Gabuzda, D. C., Pushkarev, A. B., & Garnich, N. N. 2001, MNRAS, 327, 1
- Garrett, M. A. 2005, in EAS Publications Series, Vol. 15, EAS Publications Series, ed. L. I. Gurvits, S. Frey, & S. Rawlings, 73-91
- Goddi, C., Greenhill, L. J., Chandler, C. J., et al. 2009, ApJ, 698, 1165

- Gómez, J. L., Alberdi, A., & Marcaide, J. M. 1993, A&A, 274, 55 Gómez, J. L., Lobanov, A. P., Bruni, G., et al. 2016, ApJ, 817, 96 Gómez, J. L., Marscher, A. P., Alberdi, A., Jorstad, S. G., & García-Miró, C. 2000, Science, 289, 2317
- Gómez, J. L., Marscher, A. P., Jorstad, S. G., Agudo, I., & Roca-Sogorb, M. 2008, ApJ, 681, L69
- Gómez, J. L., Marti, J. M., Marscher, A. P., & Ibanez, J. M. 1997a, Vistas in Astronomy, 41, 79
- Gómez, J. L., Martí, J. M., Marscher, A. P., Ibáñez, J. M., & Alberdi, A. 1997b, ApJ, 482, L33
- Gómez, J. L., Marti, J. M. A., Marscher, A. P., Ibanez, J. M. A., & Marcaide, J. M. 1995, ApJL, 449, L19 Gray, M. D., Wittkowski, M., Scholz, M., et al. 2009, MNRAS, 394, 51
- Greenhill, L. J., Goddi, C., Chandler, C. J., Matthews, L. D., & Humphreys, E. M. L. 2013, ApJL, 770, L32
- Hada, K., Doi, A., Kino, M., et al. 2011, Nature, 477, 185
- Han, S.-T., Lee, J.-W., Kang, J., et al. 2013, PASP, 125, 539
- Helmboldt, J. F., Taylor, G. B., Tremblay, S., et al. 2007, ApJ, 658, 203 Hirota, T., Tsuboi, M., Kurono, Y., et al. 2014, PASJ, 66, 106
- Höfner, S., Gautschy-Loidl, R., Aringer, B., & Jørgensen, U. G. 2003, A&A, 399, 589
- Hovatta, T., Aller, M. F., Aller, H. D., et al. 2014, AJ, 147, 143
- Hovatta, T., Lister, M. L., Aller, M. F., et al. 2012, AJ, 144, 105
- Humphreys, E. M. L., Gray, M. D., Yates, J. A., et al. 2002, A&A, 386, 256
- Imai, H., Nakashima, J.-I., Deguchi, S., et al. 2010, PASJ, 62, 431
- Johnson, M. D., Fish, V. L., Doeleman, S. S., et al. 2015, Science, 350, 1242 Jorstad, S. G., Marscher, A. P., Smith, P. S., et al. 2013, ApJ, 773, 147
- Jung, T., Dodson, R., Han, S.-T., et al. 2015, Journal of Korean Astronomical Society, 48, 277
- Jung, T., Sohn, B. W., & Byun, D.-Y. 2012, in Proceedings of the 11th European VLBI Network Symposium & Users Meeting. 9-12 October, 2012. Bordeaux (France). Online at http://pos. sissa. it/cgi-bin/reader/conf. cgi? confid= 178, id. 60, 60 Jung, T., Sohn, B. W., Kobayashi, H., et al. 2011, PASJ, 63, 375
- Kalamkar, M., Casella, P., Uttley, P., et al. 2016, MNRAS, 460, 3284
- Kamohara, R., Bujarrabal, V., Honma, M., et al. 2010, A&A, 510, A69
- Kim, H.-G., Han, S.-T., Sohn, B. W., et al. 2004, in European VLBI Network on New Developments in VLBI Science and Technology, ed. R. Bachiller, F. Colomer, J.-F. Desmurs, & P. de Vicente, 281-284
- Königl, A. 1981, ApJ, 243, 700
- Kovalev, Y. Y., Kellermann, K. I., Lister, M. L., et al. 2005, AJ, 130, 2473
- Kovalev, Y. Y., Lobanov, A. P., Pushkarev, A. B., & Zensus, J. A. 2008, A&A, 483, 759
- Kovalev, Y. Y., Petrov, L., Fomalont, E. B., & Gordon, D. 2007, AJ, 133, 1236
- Krichbaum, T. P., Roy, A., Lu, R.-S., et al. 2014, in Proceedings of the 12th European VLBI Network Symposium and Users Meeting (EVN 2014). 7-10 October 2014. Cagliari, Italy. Online at http://pos.sissa.it/cgi-bin/reader/conf.cgi?confid=230,id.13, 13 Krumholz, M. R., Klein, R. I., McKee, C. F., Offner, S. S. R., & Cunningham, A. J. 2009, Science, 323, 754

- Kuiper, R., Klahr, H., Beuther, H., & Henning, T. 2010, ApJ, 722, 1556
- Laing, R. A. 1981, ApJ, 248, 87
- Lanyi, G. E., Boboltz, D. A., Charlot, P., et al. 2010, AJ, 139, 1695
- Lee, S.-S., Lobanov, A. P., Krichbaum, T. P., et al. 2008, AJ, 136, 159
- Lee, S.-S., Petrov, L., Byun, D.-Y., et al. 2014, AJ, 147, 77
- Lister, M. L. & Homan, D. C. 2005, AJ, 130, 1389
- Lobanov, A. P. 1998, A&A, 330, 79
- Lockett, P. & Elitzur, M. 1992, ApJ, 399, 704
- Marscher, A. P., Jorstad, S. G., D'Arcangelo, F. D., et al. 2008, Nature, 452, 966
- Marscher, A. P., Jorstad, S. G., Gómez, J. L., et al. 2002, Nature, 417, 625
- Marscher, A. P., Jorstad, S. G., Larionov, V. M., et al. 2010, ApJ, 710, L126
- Marti-Vidal, I., Abellan, F. J., Marcaide, J. M., et al. 2016, ArXiv e-prints [arXiv:arXiv:1607.05089]
- Matthews, L. & Crew, G. 2015, Summary of the First ALMA Phasing Project (APP) Commissioning and Science Verification Mission: 2015 January 6-13, Tech. rep., NRAO
- Matthews, L. D., Greenhill, L. J., Goddi, C., et al. 2010, ApJ, 708, 80 McKinney, J. C., Tchekhovskoy, A., & Blandford, R. D. 2013, Science, 339, 49
- Middelberg, E., Roy, A. L., Walker, R. C., & Falcke, H. 2005, A&A, 433, 897
- Miller-Jones, J. C. A., Fender, R. P., & Nakar, E. 2006, MNRAS, 367, 1432
- Miller-Jones, J. C. A., Rupen, M. P., Türler, M., et al. 2009, MNRAS, 394, 309
- Mimica, P., Aloy, M.-A., Agudo, I., et al. 2009, ApJ, 696, 1142 Mirabel, I. F. & Rodríguez, L. F. 1994, Nature, 371, 46

- Mirabel, I. F. & Rodriguez, L. F. 1999, Annu. Rev. Astro. Astrophys., 37, 409
- Miyoshi, M., Matsumoto, K., Kameno, S., Takaba, H., & Lwata, T. 1994, Nature, 371, 395
- Molina, S. N., Agudo, I., Gómez, J. L., et al. 2014, A&A, 566, A26
- Moscadelli, L., Cesaroni, R., Rioja, M. J., Dodson, R., & Reid, M. J. 2011, A&A, 526, A66
- Neufeld, D. A. & Melnick, G. J. 1991, ApJ, 368, 215

- Neufeld, D. A., Wu, Y., Kraus, A., et al. 2013, ApJ, 769, 48
- Oh, S.-J., Roh, D.-G., Wajima, K., et al. 2011, PASJ, 63, 1229
- Ojha, R., Fey, A. L., Charlot, P., et al. 2005, AJ, 130, 2529
- Ojha, R., Fey, A. L., Johnston, K. J., et al. 2004, AJ, 127, 3609
- Ojha, R., Kadler, M., Böck, M., et al. 2010, A&A, 519, A45
- Olofsson, H., Rydbeck, O. E. H., Lane, A. P., & Predmore, C. R. 1981, ApJL, 247, L81
- O'Sullivan, S. P. & Gabuzda, D. C. 2009a, MNRAS, 400, 26
- O'Sullivan, S. P. & Gabuzda, D. C. 2009b, MNRAS, 393, 429
- Perucho, M., Bosch-Ramon, V., & Khangulyan, D. 2010, A&A, 512, L4
- Petrov, L. 2016, ArXiv e-prints [arXiv:1610.04951]
- Petrov, L., Hirota, T., Honma, M., et al. 2007, AJ, 133, 2487 Petrov, L., Kovalev, Y. Y., Fomalont, E. B., & Gordon, D. 2008, AJ, 136, 580
- Petrov, L., Lee, S.-S., Kim, J., et al. 2012, AJ, 144, 150 Pollack, L. K., Taylor, G. B., & Zavala, R. T. 2003, ApJ, 589, 733
- Porcas, R. W. & Rioja, M. J. 2002, in Proceedings of the 6th EVN Symposium, ed. E. Ros, R. W. Porcas, A. P. Lobanov, & J. A. Zensus, 65
- Porth, O., Fendt, C., Meliani, Z., & Vaidya, B. 2011, ApJ, 737, 42
- Pudritz, R. E. & Banerjee, R. 2005, in IAU Symposium, Vol. 227, Massive Star Birth: A Crossroads of Astrophysics, ed. R. Cesaroni, M. Felli, E. Churchwell, & M. Walmsley, 163-173
- Reid, M. J. & Honma, M. 2014, ARAA, 52, 339
- Reid, M. J., Menten, K. M., Greenhill, L. J., & Chandler, C. J. 2007, ApJ, 664, 950
- Richards, A. M. S., Etoka, S., Gray, M. D., et al. 2012, A&A, 546, A16
- Rioja, M. & Dodson, R. 2011, AJ, 141, 114
- Rioja, M., Dodson, R., Malarecki, J., & Asaki, Y. 2011, AJ, 142, 157
- Rioja, M. J., Dodson, R., Jung, T., & Sohn, B. W. 2015, AJ, 150, 202
- Rioja, M. J., Dodson, R., Jung, T., et al. 2014, AJ, 148, 84
- Rioja, M. J., Dodson, R., Kamohara, R., et al. 2008, PASJ, 60, 1031 Sánchez Contreras, C., Desmurs, J. F., Bujarrabal, V., Alcolea, J., & Colomer, F. 2002, A&A, 385, L1
- Shen, Z.-Q., Lo, K. Y., Liang, M.-C., Ho, P. T. P., & Zhao, J.-H. 2005, Nature, 438, 62
- Shu, F. H. 1985, in IAU Symposium, Vol. 106, The Milky Way Galaxy, ed. H. van Woerden, R. J. Allen, & W. B. Burton, 561–566 Shu, F. H., Najita, J., Ostriker, E. C., & Shang, H. 1995, ApJL, 455, L155
- Soker, N. 2001, ApJ, 558, 157
- Sokolovsky, K. V., Kovalev, Y. Y., Pushkarev, A. B., & Lobanov, A. P. 2011, A&A, 532, 38
- Soria-Ruiz, R., Alcolea, J., Colomer, F., et al. 2004, A&A, 426, 131
- Stirling, A. M., Spencer, R. E., de la Force, C. J., et al. 2001, MNRAS, 327, 1273
- Taylor, G. B., Vermeulen, R. C., Readhead, A. C. S., et al. 1996, ApJS, 107, 37
- Tetarenko, A., Sivakoff, G. R., Young, K., Wouterloot, J. G. A., & Miller-Jones, J. C. 2015, The Astronomer's Telegram, 7708
- Tingay, S. J., Jauncey, D. L., Preston, R. A., et al. 1995, Nature, 374, 141
- Vaidya, B. & Goddi, C. 2013, MNRAS, 429, L50
- van Leeuwen, F., ed. 2007, Astrophysics and Space Science Library, Vol. 350, Hipparcos, the New Reduction of the Raw Data
- Walker, R. C., Dhawan, V., Romney, J. D., Kellermann, K. I., & Vermeulen, R. C. 2000, ApJ, 530, 233
- Wolfire, M. G. & Cassinelli, J. P. 1987, ApJ, 319, 850
- Yun, Y., Cho, S.-H., Imai, H., et al. 2016, ApJ, 822, 3
- Zamaninasab, M., Clausen-Brown, E., Savolainen, T., & Tchekhovskoy, A. 2014, Nature, 510, 126
- Zavala, R. T. & Taylor, G. B. 2005, ApJ, 626, L73